3 (Sem-6) CHM M 4

On which facto 0202e biological system

the Hill constant depends? YRTSIMEND

(d) Which comple (noise is used for the

treatment of heavy marked toxicity?
Paper: 6.4

(Inorganic Chemistry)

Full Marks: 60

Time : Three hours

The figures in the margin indicate full marks for the questions.

1. Answer the following:

 $1 \times 7 = 7$

2. (a) In the reaction

(a) What states are expected in an octahedral field from a ¹G free ion term?

- (b) Between [Ni(CN)₄]⁻² and [Co(NH₃)₆]⁺³ complex ions, which one is kinetically inert?
 - (c) On which factor of the biological system the Hill constant depends?
 - (d) Which complex of EDTA is used for the treatment of heavy metal toxicity?
 - (e) Name the actinide whose compound is used as a reagent in your laboratory.
 - (f) Why are lanthanides known as inner transition elements?
 - (g) Complete the following:

$$^{30}_{15}P \longrightarrow ^{30}_{14}Si + \dots$$

2. (a) In the reaction

$$[Co(NH_3)_4Cl_2]^+ + Cl^- \longrightarrow [Co(NH_3)_3Cl_3] + NH_3$$

only one isomer of the complex product
is obtained. Is the initial complex *cis*
or *trans*?

3 (Sem-6) CHM M4/G 2

(b) Define the following terms:

effect among the essential trace

reaction? How is it calculated?

- more intense d rotqoos (i)
- (c) Write briefly about the antagonistic
- (c) Describe the use of gold compounds in medicine.
- Calculate the binding energy of an α -particle. Mass of proton = 1.0078 amu, mass of neutron = 1.0089 amu, mass of α -particle = 4.0084 amu. Express the result in MeV.
- 3. Answer any three questions: 5×3=15
- (a) How three isomeric $[Pt(NH_3)(Py)ClBr]$ complexes can be synthesized from $[PtCl_4]^{-2}$ ion?

- Explain which one of the following complexes should have comparatively more intense d-d transition: Ni(CO)4 or Fe(CO)5
- Write briefly about the antagonistic effect among the essential trace clements? lo scurbe the use of stnemele
- (d) What is meant by Q-value of nuclear reaction? How is it calculated? 7=8+2) Calculate the binding energy of an

- Write the outer electronic configuration of promethium. Why is it not found in nature? ? + 0084 arr
- Answer any three of the following: 10×3=30
 - (a) With the help of Orgel diagram. explain how many absorption bands are expected the electronic spectrum of $[Cr(H_2O)_6]^{+3}$ ion. Level the energy levels and assign the transitions. 3+(1+1)=5

- Discuss the theory of calorimetric determination of metals.
- Write about the absorption and (b) (i) metabolic function of vitamin B12.

5

- (ii) How does cyanide ion act as a poison in the human body? What is the antidote of cyanide poisoning? 3+2=5
- Write about the radioactive disintegration series.
 - Write the theory of redox titration.

(d) (i) s What are complexones? Name the simplest complexone with its formula. Which complexone is widely used in analytical chemistry and why? How the charge of cation, pH of solution and stability of the complexes formed with it are correlated? 1/2+1/2+3+3=7

2

rimetric	nlen 1	(ii) Dientse the theory of
5	(ii)	The reaction of actinides with hot
	more	water is faster than expected.
		Why? white about (i) (d) 3
	vitan	metabolic function of
(e)	(i)	Write about the characteristics of
		'London smog' and 'Los Angeles
		What is the 'gome'
3+2=5		poisoning?
oactive	(ii)	Explain the mechanism of Na+-K+
7		5 disintegration.qmuq
itration.	dox t	(ii) Write the theory of re
E (f)	(i)	The absorption band of Ce (III) ion
ame the	SPN	is broad, while that of other
rith its	ne w	lanthanide ions are sharp. Explain.
gi sone is	inple	formula. Which con
		widely used in analyti
		Actinides have a greater
with 12 1+3+3=7	rmed	tendency to form complexes than
0.64		lanthanides. Explain. 4

6

(iii) A piece of wood taken from a cave is found to have C-14 activity only 0.636 times that of fresh wood sample. Estimate the age of the wood sample. The half-life of C-14 is 5700 years.